



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

303. Proposed by FRANCIS RUST, C. E., Allegheny, Pa.

Prove that the pedal line of any point on a triangle's circum-circle bisects the distance from this point to the triangle's ortho-center.

304. Proposed by G. W. GREENWOOD, M. A., Dunbar, Pa.

Find the tangent at the points $(a, 0)$ and $(0, a)$ to the locus $x^3 + y^3 = a^3$, and show that these points are points of inflection.

305. Proposed by J. J. QUINN, Ph. D., Scottdale, Pa.

1. Suppose two radii R and R_1 revolve uniformly in the ratio 2 : 3. Find the equation of the locus of the intersection of R with the chord drawn from the end of the diameter to the extremity of R_1 . 2. If the chord be drawn to the end of the diameter to the extremity of R , what is the locus of the intersection with R_1 ? 3. Show how an angle can be trisected by means of this curve.

CALCULUS.

230. Proposed by C. N. SCHMALL, College of the City of New York.

The greatest rectangle is inscribed in an ellipse, and the greatest ellipse in that rectangle, again the greatest rectangle in that (second) ellipse, and the greatest ellipse in that (second) rectangle, and so on *ad infinitum*; show that the sum of all the inscribed rectangles is equal to the area of the rectangle circumscribed about the given ellipse.

231. Proposed by EVA S. MAGLOTT, A. M., Professor of Mathematics, Ohio Northern University, Ada, O.

If a right circular cone stands on an ellipse, prove that the convex surface of the cone is $\frac{1}{2}\pi(OA + OA')(OA.OA')^{\frac{1}{2}} \sin \alpha$, where O is the vertex of the cone, A and A' the extremities of the major axis of the ellipse, and α is the semi-angle of the cone at the vertex, using the formula $ds = \frac{1}{2}\rho\sqrt{(\rho^2 + p^2)}d\theta$, where p is the perpendicular from the vertex to the base of the cone, ρ the distance from the foot of the perpendicular to any point in the perimeter of the base, and θ the angle between the major axis and ρ .

MECHANICS.

194. Proposed by W. J. GREENSTREET, M. A., Editor of the Mathematical Gazette, Stroud, England.

A body has a plane face resting on a rough wedge. The wedge is on a rough inclined plane, thick end down and thin edge horizontal. Find the condition that the body will slide down the wedge with constant acceleration, the wedge not slipping the while. Discuss the case in which the angle of friction for wedge and plane is greater than the angle of inclination of the plane.